311 within these 800 acres, much is occupied by the streams, calculated at their
241 flood maxima which define non-gardenable areas, while other parts are inadequate in soils or water for use as gardens. Consequently, the actually tillable areas comprise a maximum of about 200 acres or about 0.5% of the land surface. It should be noted, of course, that all these 200 acres are not usable for crops since those in the *montes* cannot be used for cane and bananas, while the wet areas nearest the streams cannot be used for corn or manioc. Thus the maximum area for the staple crops, manioc, corn, and bananas, is about 145 acres (see Table I)[A]. [11A]

168 Using Carneiro's equations, [12A] we may estimate the maximum population
311 which would be in equilibrium with this acreage, estimating .5 acres as an approximate amount of land required to provide the average person with the food he ordinarily derives from cultivated plants per year. [13A] The calculation gives us 92 persons (see Figure 1). [14A] We may also estimate the number of years (L) the group may stay in the present locality with its present population of 24 before wearing out the available tillable land so that they would have to move (see Figure 2). This would be 50 years. Since the entire fallow and planting cycle of any piece of land is only 13 years, under present circumstances, the Totúdoro Yaruro will not have to move from their present territory at all, in view of their small and decreasing population.

[128]

Figure 1

$$P = \frac{\frac{T}{(R+Y)} \times Y}{A}$$

$$P = \frac{\frac{200}{10+3} \times 3}{.5}$$

P = 92 (=67 if R is 15)

Figure 2

$$L = \frac{\frac{T}{P \times A}}{Y}$$

$$L = \frac{\frac{200}{24 \times .5}}{3}$$

L = 50 years

R = year of fallow
Y = years of productivity
A = land used to provide plant food for one person a year
P = population
L = number of years group located in one place
T = acres of cultivable land within walking-working distance of village

[262] The ideal figures given above would be somewhat modified upwards by the additional foods provided by fishing, hunting, and gathering which, in the wet season, constitute about 30% of the diet and, in the dry, perhaps

[A] [for Table I see pp. 28-30, category 116--Analyst]
[11,12,13,14] [for footnotes, see p. 43, category 116--Analyst]

50% of the diet or a little less. The ideal figures would again be reduced, on the other hand, by allowing for the predations on gardens of animals, especially the pigs, which eat manioc and corn, the Guahibo Indians who allegedly steal sugar cane, and insect losses. Our figures, then, will be modified only a few units by the additional sources of foods. We may thus revise our population figure to between 100 and 105 (or, if a 15-year fallow period is assumed, the figure would be 75-80), figures which correspond to the Yaruro's own ideal of what a proper population for this area should be. If the population increased beyond this number of people, the Yaruro would have to exploit their gardens at a rate greater than the rate of land reconstitution, and also expand hunting, fishing, and gathering beyond the average long-term carrying capacity of the savanas and the rivers, thus reducing their potential productivity. Adding together the maximum populations supportable by these food resources, as well as by the recently introduced pigs, a figure approximating 125-150 persons might be reasonable for this particular territory. This gives a density of 2 persons per sq. mile, a characteristic figure for tropical forest tribes.[15]

If the population rose beyond this maximum capacity of the land-technology relationship, groups would either have to fission off, go on the warpath, regulate population by any of a number of internal institutions such as infanticide, or die either from extreme famines or from steady nutritional deficiency. Under present population conditions none of these is necessary, yet both fissioning and population regulation by contraception occur. Fissioning appears to be fostered by the search for settlement locations requiring minimum labor for maximum return and also to establish community authority under the person leading the fissioning group independent of that excercised by the community from which fissioning took place. Contraception appears to be a matter of choice revolving about maintaining a certain freedom from the cares of motherhood until such time as the marriage partners are reasonably convinced that the marriage is permanent. Indirectly, this contributes to the distribution of more effective female producers among the more effective male producers in the most productive areas, since a second marriage often involves shifting to a generally more effective husband whereas a shift to a less effective husband is scarcely to be found. Both fissioning and contraception, however, give evidence of techniques which were developed to handle man-food balance in happier days.

A number of formal characteristics of any ecology may permit us to deduce the form of society which would be expected on the basis of such characteristics. In so far as the ecological forms were to remain essentially unchanged over great periods of time, no changes in the forms of the social order attached to the ecological conditions would be expected. An internal equilibrium is maintained until such time as a dynamic imbalance occurs because of changes from internal or external sources. It is essential to examine to what extent the internal equilibrium prevents or fosters the occurence of dynamic imbalances in any particular ecological setting. We shall return to this question after the following itemization of Yaruro ecological forms.

[15] [for footnote see p. 43, category 116 -- Analyst]